

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 November 2001 (29.11.2001)

PCT

(10) International Publication Number
WO 01/90078 A1

(51) International Patent Classification⁷: **C07D 231/14**,
231/20, 231/22, A61K 31/415, A61P 1/04

(74) Agents: **FISHER, Adrian, John** et al.; Carpmals &
Ransford, 43 Bloomsbury Square, London WC1A 2RA
(GB).

(21) International Application Number: **PCT/GB01/01976**

(22) International Filing Date: **4 May 2001 (04.05.2001)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
0011095.7 **8 May 2000 (08.05.2000)** **GB**

(71) Applicant (for all designated States except US): **JAMES
BLACK FOUNDATION LIMITED** [GB/GB]; 68 Half
Moon Lane, Dulwich, London SE24 9JE (GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and




(75) Inventors/Applicants (for US only): **MCDONALD,
Iain, Mair** [GB/GB]; 68 Half Moon Lane, Dulwich,
London SE24 9JE (GB). **LOW, Caroline, Minli, Rachel**
[GB/GB]; 68 Half Moon Lane, Dulwich, London SE24
9JE (GB). **STEEL, Katherine, Isobel, Mary** [GB/GB];
68 Half Moon Lane, Dulwich, London SE24 9JE (GB).
SPENCER, John [GB/GB]; 68 Half Moon Lane, Dul-
wich, London SE24 9JE (GB).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: **PYRAZOLE DERIVATIVES AND THEIR USE AS GASTRIN AND CHOLECYSTOKININ RECEPTOR LIGANDS**



(57) Abstract: Compounds of formula (I), (IIa) or (IIb) wherein R¹-R⁶, Z, Q and n are as defined in claim 1 and their pharmaceuti-
cally acceptable salts are ligands at gastrin and/or cholecystokinin receptors. Compositions comprising a compound of formula (I),
(IIa) or (IIb) are also described.

PYRAZOLE DERIVATIVES AND THEIR USE AS GASTRIN AND CHOLECYSTOKININ IN RECEPTOR LIGANDS

This invention relates to gastrin and cholecystokinin (CCK) receptor ligands. (The receptor previously known as the CCK_B/gastrin receptor is now termed the CCK₂ receptor). The invention also relates to methods for preparing such ligands and to compounds which are useful intermediates in such methods. The invention further relates to pharmaceutical compositions comprising such ligands and methods for preparing such pharmaceutical compositions.

- 10 Gastrin and the cholecystokinins are structurally related neuropeptides which exist in gastrointestinal tissue and the central nervous system (Mutt V., *Gastrointestinal Hormones*, Glass G.B.J., ed., Raven Press, New York, p. 169; Nissson G., *ibid.*, p. 127).

Gastrin is one of the three primary stimulants of gastric acid secretion. Several forms of gastrin are found including 34-, 17- and 14-amino acid species with the minimum active fragment being the C-terminal tetrapeptide (TrpMetAspPhe-NH₂) which is reported in the literature to have full pharmacological activity (Tracey H.J. and Gregory R.A., *Nature* (London), 1964, **204**, 935). Much effort has been devoted to the synthesis of analogues of this tetrapeptide (and the N-protected derivative Boc-TrpMetAspPhe-NH₂) in an attempt to elucidate the relationship between structure and activity.

Natural cholecystokinin is a 33 amino acid peptide (CCK-33), the C-terminal 5 amino acids of which are identical to those of gastrin. Also found naturally is the C-terminal octapeptide (CCK-8) of CCK-33.

25

The cholecystokinins are reported to be important in the regulation of appetite. They stimulate intestinal mobility, gall bladder contraction, pancreatic enzyme secretion and are known to have a trophic action on the pancreas. They also inhibit gastric emptying and have various effects in the central nervous system.

30

Compounds which bind to cholecystokinin and/or gastrin receptors are important because of their potential pharmaceutical use as antagonists or partial agonists of the natural peptides.


- 5 A number of gastrin antagonists have been proposed for various therapeutic applications, including the prevention of gastrin-related disorders, gastrointestinal ulcers, Zollinger-Ellison syndrome, antral G cell hyperplasia and other conditions in which lower gastrin activity or lower acid secretion is desirable. The hormone has also been shown to have a trophic action on cells and so an antagonist may be expected to be useful in the treatment
10 of cancers, particularly in the stomach and the colon.

Possible therapeutic uses for cholecystokinin antagonists include the control of appetite disorders such as anorexia nervosa and the treatment of pancreatic inflammation, biliary tract disease and various psychiatric disorders. Other possible uses are in the potentiation
15 of opiate (for example morphine) analgesia and in the treatment of cancers, especially of the pancreas. Moreover, ligands for cholecystokinin receptors in the brain (so-called CCK₂ receptors) have been claimed to possess anxiolytic activity.

PCT/GB99/03733 describes a class of compounds having gastrin antagonist activity. This
20 class of compounds is typically characterised by a 5-membered ring, preferably an imidazole or pyrrole, having two hydrocarbyl substituents and an amide or urea-type substituent.

According to the present invention, there are provided compounds of formula (I), (IIa) or
25 (IIb).

3



wherein

n is from 1 to 4;

R^1 is H or C_1 to C_{15} hydrocarbyl wherein up to three C atoms may optionally be replaced by N, O and/or S atoms and up to three H atoms may optionally be replaced by halogen atoms;

R^2 is selected from H, Me, Et, Pr and OH, each R^2 being independently selected from H, Me, Et, Pr and OH when n is greater than 1;

R^3 (when n is 1) is selected from H, Me, Et and Pr; or (when n is greater than 1) each R^3 is independently selected from H, Me, Et and Pr, or two R^3 groups on neighbouring carbon atoms are linked to form a C_3 to C_6 carbocyclic ring, or two R^3 groups are absent from neighbouring carbon atoms which are linked by a double bond; or R^2 and R^3 on the same carbon atom together represent an =O group;

R^4 is H or C_1 to C_{15} hydrocarbyl wherein up to two C atoms may optionally be replaced by N, O and/or S atoms and up to three H atoms may optionally be replaced by halogen atoms;


R^5 is H, OH or C_1 to C_{15} hydrocarbyl wherein up to two C atoms may optionally be replaced by N, O and/or S atoms and up to three H atoms may optionally be replaced by halogen atoms;

R^6 is H or C_1 to C_{15} hydrocarbyl wherein up to two C atoms may optionally be replaced by N, O and/or S atoms and up to three H atoms may optionally be replaced by halogen atoms;


Z is $-(NR^7)_a-CO-(NR^8)_b-$ (wherein a is 0 or 1, b is 0 or 1, and R^7 and R^8 are independently selected from H, Me, Et, Pr or Bn),

$-CO-NR^7-CH_2-CO-NR^8-$, $-CO-O-$, $-CH_2-CH_2-$, $-CH=CH-$, $-CH_2-NR^8-$ or a bond;

5 Q is $-R^9V$, or



wherein R^9 is $-CH_2-$; $-CH_2-CH_2-$; or



10 or R^9 and R^8 , together with the nitrogen atom to which R^8 is attached, form a piperidine or pyrrolidine ring which is substituted by V;

V is $-CO-NH-SO_2-Ph$, $-SO_2-NH-CO-Ph$, $-CH_2OH$, or a group of the formula $-R^{10}U$, (wherein U is $-COOH$, tetrazolyl, $-CONHOH-$ or $-SO_3H$; and R^{10} is a bond; C_1 to C_6 hydrocarbylene, optionally substituted by hydroxy, amino or acetamido; $-O-(C_1$ to C_3 alkylene)-; $-SO_2NR^{11}-CHR^{12}-$;

15 $-CO-NR^{11}-CHR^{12}-$, R^{11} and R^{12} being independently selected from H and methyl; or $-NH-(CO)_c-CH_2-$, c being 0 or 1);

T is C_1 to C_6 hydrocarbyl, $-NR^7R^8$ (wherein R^7 and R^8 are as defined above),

$-OMe$, $-OH$, $-CH_2OH$, halogen or trihalomethyl;

m is 1 or 2;


20 p is from 0 to 3; and

q is from 0 to 2, with the proviso that q is 1 or 2 when Z is a bond);

or a pharmaceutically acceptable salt thereof.

In certain compounds according to the invention, Z is $-(NR^7)_a-CO-(NR^8)_b-$, $-CO-NH-$
 25 $CH_2-CO-NH-$ or a bond; Q is

5




V is $-\text{CO}-\text{NH}-\text{SO}_2-\text{Ph}$, $-\text{SO}_2-\text{NH}-\text{CO}-\text{Ph}$, $-\text{OCH}_2\text{COOH}$, tetrazolyl or $-(\text{CH}_2)_s\text{COOH}$, wherein s is from 0 to 2; and T is C_1 to C_6 hydrocarbyl, $-\text{NR}^7\text{R}^8$, $-\text{OMe}$, $-\text{OH}$, $-\text{CH}_2\text{OH}$ or halogen

5

A further group of compounds according to the invention are those in Z is $-(\text{NR}^7)_a-\text{CO}-(\text{NR}^8)_b-$, Q is $-(\text{CH}_2)_r\text{COOH}$, wherein r is from 1 to 3; and T is C_1 to C_6 hydrocarbyl, $-\text{NR}^7\text{R}^8$, $-\text{OMe}$, $-\text{OH}$, $-\text{CH}_2\text{OH}$ or halogen.

10 A still further group of compounds according to the invention are those in which $-\text{Z}-\text{Q}$ is



Preferably R^1 is C_1 to C_{12} hydrocarbyl wherein one C atom may optionally be replaced by N or O and up to three H atoms may optionally be replaced by F, Cl or Br. More preferably R^1 is H; C_3 to C_{12} alicyclic; phenyl, pyridyl, phenyl C_1 - C_3 alkyl or pyridyl C_1 - C_3 alkyl (all optionally substituted with OMe, NMe₂, CF₃, Me, F, Cl, Br or I); or C_1 to C_8 alkyl. Alicyclic groups include C_5 to C_8 cycloalkyl, C_7 to C_{10} polycycloalkyl, C_5 to C_8 cycloalkenyl and C_7 to C_{10} polycycloalkenyl, all optionally substituted with methyl. Phenyl C_1 - C_3 alkyl includes, for example, benzyl.

20 C_1 - C_3 alkyl includes, for example, benzyl.

Preferably R^2 is H, R^3 is H and n is 1, 2 or 3.

In certain compounds according to this invention R^4 is C_3 to C_{15} carbocyclic, optionally substituted with 1, 2 or 3 halogen atoms. In other compounds according to this invention, R^4 is $-\text{NH}-\text{R}^{13}$ or $-\text{OR}^{13}$, wherein R^{13} is C_3 to C_{12} carbocyclic, optionally substituted with 1, 2 or 3 halogen atoms. Preferably R^4 is $-\text{O}-\text{adamantyl}$, $-\text{O}-\text{cycloheptyl}$, $-\text{O}-\text{cyclohexyl}$ or $-\text{O}-\text{phenyl}$. Most preferably R^4 is $-\text{O}-\text{adamantyl}$.

25

Preferably R^5 is H, OH or C_1 to C_{12} hydrocarbonyl wherein one C atom may optionally be replaced by N or O and up to three H atoms may optionally be replaced by F, Cl, or Br. More preferably R^5 is H; OH; C_3 to C_{12} alicyclic; $O-C_3$ to C_{12} alicyclic; phenyl, pyridyl, phenyl C_1-C_3 alkyl, phenyl C_1-C_3 alkoxy, pyridyl C_1-C_3 alkyl, or pyridyl C_1-C_3 alkoxy (all optionally substituted with OMe, NMe₂, CF₃, Me, F, Cl, Br or I); C_1 to C_8 alkyl; or C_1 to C_8 alkoxy. Alicyclic groups include C_5 to C_8 cycloalkyl, C_7 to C_{10} polycycloalkyl, C_5 to C_8 cycloalkenyl and C_7 to C_{10} polycycloalkenyl, all optionally substituted with methyl. Phenyl C_1-C_3 alkyl includes, for example, benzyl.


10

Preferably R^6 is H or C_1 to C_{12} hydrocarbonyl wherein one C atom may optionally be replaced by N or O and up to three H atoms may optionally be replaced by F, Cl, or Br. More preferably R^6 is H; C_3 to C_{12} alicyclic; phenyl, pyridyl, phenyl C_1-C_3 alkyl or pyridyl C_1-C_3 alkyl (all optionally substituted with OMe, NMe₂, CF₃, Me, F, Cl, Br or I); or C_1 to C_8 alkyl. Alicyclic groups include C_5 to C_8 cycloalkyl, C_7 to C_{10} polycycloalkyl, C_5 to C_8 cycloalkenyl and C_7 to C_{10} polycycloalkenyl, all optionally substituted with methyl. Phenyl C_1-C_3 alkyl includes, for example, benzyl.


20

Preferably Z is $-CO-NH-$.

Preferably Q is



and more preferably



25

p is preferably 0 or 1, and q is preferably 0. If p is greater than 0, then T is preferably C_1 to C_6 hydrocarbonyl or halo.

m is preferably 1, and V is preferably $-\text{CO}_2\text{H}$, $-\text{CH}_2\text{CO}_2\text{H}$ or tetrazolyl.

R^{10} is preferably a bond, C_1 or C_2 alkylene (optionally substituted by hydroxy, amino or acetamido), $-\text{O}-(\text{C}_1 \text{ to } \text{C}_3 \text{ alkylene})-$; $-\text{SO}_2\text{NR}^{11}-\text{CHR}^{12}-$; $-\text{CO}-\text{NR}^{11}-\text{CHR}^{12}-$,

5 $-\text{NH}-(\text{CO})_c-\text{CH}_2-$, or a group of the formula




Certain compounds of the invention exist in various regioisomeric, enantiomeric, tautomeric and diastereomeric forms. It will be understood that the invention comprehends
10 the different regioisomers, enantiomers, tautomers and diastereomers in isolations from each other as well as mixtures.


In particular, the skilled person will understand that the compound of formula (IIa) wherein R^6 is H and the compound of formula (IIb) are merely tautomers of the compound of
15 formula (I) wherein R^5 is OH. These tautomers may exist in isolation from each other or as mixtures.

Compounds of present invention wherein

- (i) R^5 is H
20 (ii) $-\text{Z}-\text{Q}$ is



may be prepared by the route exemplified in Reaction Scheme 1.




Reaction Scheme 1

β -Ketoester (III) is reacted with DMF dimethylacetal to yield α,β -unsaturated ketone (IV).


- 5 Compound (IV) is treated with R^1 -NHNH₂ to furnish the corresponding hydrazone which, in the presence of base (e.g. triethylamine), cyclises to form pyrazole (V). Conversion of the ester to an acid, followed by coupling (under acid activating conditions e.g via the acid chloride using oxalyl chloride) with an ester-substituted aniline and final unmasking of this ester furnishes the target carboxylic acid (VI). When, for example, the ester(s) are benzyl
- 10 esters, the corresponding carboxylic acid(s) can be unmasked by catalytic hydrogenation. It will be readily apparent to the skilled person that other compounds of the present invention wherein R^5 is H can be prepared by appropriate manipulation of compound (V).

Compounds of the present invention wherein

- 15 (i) R^5 is C₃ to C₁₂ alicyclic; phenyl (optionally substituted with OMe, NMe₂, CF₃, Me, F, Cl, Br, I or OH); or C₁ to C₈ alkyl
- (ii) -Z-Q is



may be prepared by the route exemplified in Reaction Scheme 2.




Reaction Scheme 2

- 5 β -Ketoester (III) is reacted with R^5 -CHO/ethylenediamine diacetate to yield α,β -unsaturated ketone (VII). Treatment of ketone (VII) with R^1 -NHNH₂ affords a hydrazone which, in the presence of base (e.g. DBU); cyclises to furnish pyrazole (VIII). Pyrazole (VIII) is converted to the target carboxylic acid (IX) by the same route shown and described in Reaction Scheme 1. It will be readily apparent to the skilled person that other
- 10 compounds of the present invention wherein R^5 is C_3 to C_{12} alicyclic; phenyl (optionally substituted with OMe, NMe₂, CF₃, Me, F, Cl, Br, I or OH); or C_1 to C_8 alkyl can be prepared by appropriate manipulation of compound (VIII).


Compounds of the present invention wherein

- 15 (i) R^5 is OH
(ii) $-Z-Q$ is



may be prepared by the route exemplified in Reaction Scheme 3.

10



Reaction Scheme 3

5

β -Ketoester (III) is reacted with R^1-NHNH_2 to form a hydrazone which, in the presence of base (e.g. DBU), cyclises to form hydroxy-substituted pyrazole (X). Deprotonation of pyrazole (X) with for example NaH, followed by treatment of the resultant anion with an appropriate isocyanate (e.g. ethylcarbonylphenylisocyanate) affords an amide. Final

10 deprotection of the ester group furnishes the requisite hydroxy-substituted pyrazole (XI) and/or its tautomers (XII) and (XIII). The ratio of (XI):(XII):(XIII) will vary from 100:0:0 to 0:100:0 to 0:0:100, depending on the other substituents present on the pyrazole ring. It will be readily apparent to the skilled person that other compounds of the present invention wherein R^5 is OH can be prepared by appropriate manipulation of compound (X).

15

Still referring to Reaction Scheme 3, compounds of the present invention wherein R^5 is OMe may be prepared by alkylation of (XI) with dimethyl sulfate. It may also be prepared by alkylation of a protected precursor of (XI), with, for example, dimethyl sulfate as shown in Reaction Scheme 4.

5


Still referring to Reaction Scheme 3, compounds of the present invention wherein R^6 is Me may be prepared by alkylation of (XII) with dimethyl sulfate. It may also be prepared by alkylation of a protected precursor of (XII), with, for example, methyl iodide as shown in Reaction Scheme 5.

10


Compounds of the present invention wherein

(i) R^5 is OMe

(ii) $-Z-Q$ is



15 may be prepared by the route exemplified in Reaction Scheme 4.




Reaction Scheme 4

Hydroxypyrazole (X) is treated with base then dimethyl sulfate to afford methoxypyrazole (XIV). Bromination with, for example, NBS gives the bromo-substituted pyrazole (XV). Finally, metallation, reaction with an appropriate isocyanate (e.g. ethylcarbonylphenylisocyanate) and deprotection furnishes the requisite methoxy-substituted pyrazole compound (XVI).


Compounds of the present invention wherein

- (i) R^6 is Me
- (ii) $-Z-Q$ is



10

may be prepared by the route exemplified in Reaction Scheme 5.



Reaction Scheme 5


15

Bromination of hydroxypyrazole (X) followed by alkylation of the pyrazole nitrogen affords (XVIII). Metallation, reaction with isocyanate and deprotection as described previously furnishes the requisite pyrazole compound (XIX).

Hence, the present invention provides methods of making compounds according to formulae (I), (IIa) or (IIb).

The invention also comprehends derivative compounds ("pro-drugs") which are degraded *in vivo* to yield the species of formula (I) or (II). Pro-drugs are usually (but not always) of lower potency at the target receptor than the species to which they are degraded. Pro-drugs are particularly useful when the desired species has chemical or physical properties which make its administration difficult or inefficient. For example, the desired species may be only poorly soluble, it may be poorly transported across the mucosal epithelium, or it may have an undesirably short plasma half-life. Further discussion of pro-drugs may be found in Stella, V. J. *et al.*, "Prodrugs", *Drug Delivery Systems*, 1985, pp. 112-176, and *Drugs*, 1985, 29, pp. 455-473.

Pro-drug forms of the pharmacologically-active compounds of the invention will generally be compounds according to formula (I) or (II) having an acid group which is esterified or amidated. Included in such esterified acid groups are groups of the form $-\text{COOR}^a$, wherein R^a is C_1 to C_5 alkyl, phenyl, substituted phenyl, benzyl, substituted benzyl, or one of the following:



20

Amidated acid groups include groups of the formula $-\text{CONR}^b\text{R}^c$, wherein R^b is H, C_1 to C_5 alkyl, phenyl, substituted phenyl, benzyl, or substituted benzyl, and R^c is $-\text{OH}$ or one of the groups just recited for R^b .

25 Compounds of formula (I) or (II) having an amino group may be derivatised with a ketone or an aldehyde such as formaldehyde to form a Mannich base. This will hydrolyse with first order kinetics in aqueous solution.

Another aspect of the present invention is a pharmaceutical composition comprising a compound of formula (I) or (II) substantially as described herein before with a pharmaceutically acceptable diluent or carrier.

- 5 Yet another aspect of the present invention is a method of making a pharmaceutical composition comprising a compound of formula (I) or (II) substantially as described herein before, comprising mixing said compound with a pharmaceutically acceptable diluent or carrier.
- 10 Pharmaceutically acceptable salts of the acidic or basic compounds of the invention can of course be made by conventional procedures, such as by reacting the free base or acid with at least a stoichiometric amount of the desired salt-forming acid or base.

Pharmaceutically acceptable salts of the acidic compounds of the invention include salts with
15 inorganic cations such as sodium, potassium, calcium, magnesium, and zinc, and salts with organic bases. Suitable organic bases include N-methyl-D-glucamine, arginine, benzathine, diolamine, olamine, procaine and tromethamine.

Pharmaceutically acceptable salts of the basic compounds of the invention include salts
20 derived from organic or inorganic acids. Suitable anions include acetate, adipate, besylate, bromide, camsylate, chloride, citrate, edisylate, estolate, fumarate, gluceptate, gluconate, glucuronate, hippurate, hyclate, hydrobromide, hydrochloride, iodide, isethionate, lactate, lactobionate, maleate, mesylate, methylbromide, methylsulfate, napsylate, nitrate, oleate, pamoate, phosphate, polygalacturonate, stearate, succinate, sulfate, sulfosalicylate, tannate,
25 tartrate, terephthalate, tosylate and triethiodide.

It is anticipated that the compounds of the invention can be administered by oral or parenteral routes, including intravenous, intramuscular, intraperitoneal, subcutaneous, rectal and topical administration, and inhalation.

30

For oral administration, the compounds of the invention will generally be provided in the form of tablets or capsules or as an aqueous solution or suspension.

Tablets for oral use may include the active ingredient mixed with pharmaceutically acceptable excipients such as inert diluents, disintegrating agents, binding agents, lubricating agents, sweetening agents, flavouring agents, colouring agents and
5 preservatives. Suitable inert diluents include sodium and calcium carbonate, sodium and calcium phosphate and lactose. Corn starch and alginic acid are suitable disintegrating agents. Binding agents may include starch and gelatine. The lubricating agent, if present, will generally be magnesium stearate, stearic acid or talc. If desired, the tablets may be coated with a material such as glyceryl monostearate or glyceryl distearate, to delay
10 absorption in the gastrointestinal tract.

Capsules for oral use include hard gelatine capsules in which the active ingredient is mixed with a solid diluent and soft gelatine capsules wherein the active ingredient is mixed with water or an oil such as peanut oil, liquid paraffin or olive oil.

15

For intramuscular, intraperitoneal, subcutaneous and intravenous use, the compounds of the invention will generally be provided in sterile aqueous solutions or suspensions, buffered to an appropriate pH and isotonicity. Suitable aqueous vehicles include Ringer's solution and isotonic sodium chloride. Aqueous suspensions according to the invention
20 may include suspending agents such as cellulose derivatives, sodium alginate, polyvinylpyrrolidone and gum tragacanth, and a wetting agent such as lecithin. Suitable preservatives for aqueous suspensions include ethyl and n-propyl p-hydroxybenzoate.

Effective doses of the compounds of the present invention may be ascertained by
25 conventional methods. The specific dosage level required for any particular patient will depend on a number of factors, including severity of the condition being treated, the route of administration and the weight of the patient. In general, however, it is anticipated that the daily dose (whether administered as a single dose or as divided doses) will be in the range 0.001 to 5000 mg per day, more usually from 1 to 1000 mg per day, and most usually from 10
30 to 200 mg per day. Expressed as dosage per unit body weight, a typical dose will be expected to be between 0.01 µg/kg and 50 mg/kg, especially between 10 µg/kg and 10 mg/kg, eg. between 100 µg/kg and 2 mg/kg.

In a further aspect of the present invention there are provided pharmaceutical compositions comprising a compound according to formula (I) or (II) and a proton pump inhibitor. Compositions comprising a CCK/gastrin antagonist and a proton pump inhibitor are
 5 described in International patent application WO93/12817, incorporated herein by reference.

In one aspect of the present invention the proton pump inhibitor is

omeprazole which is 5-methoxy-2-[[[4-methoxy-3,5-dimethyl-2-pyridinyl)-
 10 methyl]sulfinyl]-1H-benzimidazole;

BY308;

SK & 95601 which is 2-[[[3-chloro-4-morpholino-2-pyridyl)methyl]sulfinyl]-5-methoxy-(1H)-benzimidazole;

SK & 96067 which is 3-butyryl-4-(2-methylphenylamino)-8-methoxyquinoline;
 15 5-trifluoromethyl-2-[4-methoxy-3-methyl-2-pyridyl-methyl]-thio-[1H]-benzimidazole;

or pharmaceutically acceptable salts thereof.

These proton pump inhibitors are described and claimed in US Patents 4,472,409 and
 20 4,255,431. These patents are incorporated herein by reference.

In a further aspect of the present invention, the proton pump inhibitor is

lansoprazole which is 2-[[[3-methyl-4-(2,2,2-trifluoroethoxy)-2-pyridinyl]methyl]sulfinyl]-1H-benzimidazole;

25 pantoprazole which is 5-(difluoromethoxy)-2-[[[3,4-dimethoxy-2-pyridinyl)methyl]sulfinyl]-1H-benzimidazole;

perprazole;

rabeprazole which is 2-[[[4-(3-methoxypropoxy)-3-methylpyridin-2-yl]methylsulfinyl]-1H-benzimidazole;

30 [[4-(2,2,2-trifluoroethoxy)-3-methyl-2-pyridyl]-methyl]sulfenamide;

(Z)-5-methyl-2-[2-(1-naphthyl)ethenyl]-4-piperidinopyridine HCl;

2-(4-cyclohexyloxy-5-methylpyridin-2-yl)-3-(1-naphthyl)-1-propanol;

- methyl 2-cyano-3-(ethylthio)-3-(methylthio)-2propenoate;
 2-((4-methoxy-2-pyridyl)methylsulfinyl)-5-(1,1,2,2-tetrafluoroethoxy)-1H-benzimidazole sodium;
 2-[[[4-(2,2,3,3,4,4,4-heptafluorobutoxy)-2-pyridyl)methyl]sulfinyl]-1H-thieno [3,4-
 5 d]imidazole;
 2-[[[4-(2,2,2-trifluoroethoxy)-3-methyl-2-pyridyl)methyl]sulfinyl]-1H-benzimidazole;
 2-[[[4-(2,2,2-trifluoroethoxy)-3-methyl-2-pyridyl)methyl]sulfinyl]-1H-benzimidazole;
 10 2-methyl-8-(phenylmethoxy)-imidazo(1,2-A)-pyridine-3-acetonitrile;
 (2-((2-dimethylaminobenzyl)sulfinyl)-benzimidazole);
 4-(N-allyl-N-methylamino)-1-ethyl-8-((5-fluoro-6-methoxy-2-benzimidazolyl)sulfinylmethyl)-1-ethyl 1,2,3,4-tetrahydroquinolone;
 2-[[[2-(2-dimethylaminophenyl)methyl]sulfinyl]-4,7-dimethoxy-1H-benzimidazole;
 15 2-[[[2-(2-pyridyl)phenyl]sulfinyl]-1H-benzimidazole;
 (2-[[[2-(2-amino-4-methylbenzyl)sulfinyl]-5-methoxybenzo[d]imidazole;
 (4-(2-methylpyrrol-3-yl)-2-guanidisoithiazole);
 4-(4-(3-(imidazole)propoxy)phenyl)-2-phenylthiazole;
 (E)-2-(2-(4-(3-(dipropylamino)butoxy)phenyl)-ethenyl)benzoxazole;
 20 (E)-2-(2-(4-(3-(dipropylamino)propoxy)phenyl)ethenyl)-benzothiazole;
 Benzeneamine, 2-[[[5-methoxy-1H-benzimidazol-2-yl)sulfinyl)methyl]-4-methyl-;
 Pumilacidin A;
 2,3-dihydro-2-methoxycarbonylamino-1,2-benzisothiazol-3-one;
 2-(2-ethylaminophenylmethylsulfinyl)-5,6-dimethoxybenzimidazole;
 25 2-methyl-8-(phenylmethoxy)imidazo[1,2-a]pyridine-3-acetonitrile;
 3-amino-2-methyl-8-phenylmethoxyimidazo[1,2-a]-pyrazine HCl;
 2-[[[3-chloro-4-morpholino-2-pyridyl)methyl]-sulfinyl]-5-methoxy-(1H)-benzimidazole;
 [3-butyryl-4-(2-methylphenylamino)-8-methoxy-quinoline);
 30 2-indanyl 2-(2-pyridyl)-2-thiocarbamoylacetate HCl;
 2,3-dihydro-2-(2-pyridinyl)-thiazolo (3,2-a) - benzimidazole;
 3-cyanomethyl-2-methyl-8-(3-methyl-2-butenyloxy)- (1,2-a)imidazopyridine;

zinc L-carnosine;
or pharmaceutically acceptable salts thereof.

Rabeprazole is described in US patent 5,045,552. Lansoprazole is described in US patent
5 4,628,098. Pantoprazole is described in US patent 4,758,579. These patents are
incorporated herein by reference.

Preferably, the proton pump inhibitor is selected from (RS)-rabeprazole, (RS)-omeprazole,
lansoprazole, pantoprazole, (R)-omeprazole, (S)-omeprazole, perprazole, (R)-rabeprazole,
10 (S)-rabeprazole, or the alkaline salts thereof. The alkaline salts may be, for example, the
lithium, sodium, potassium, calcium or magnesium salts.

Compositions of this invention comprising a compound of formula (I) or (II) and a proton
pump inhibitor may be administered as described above. Preferably the dose of each of the
15 active ingredients in these compositions will be equal to or less than that which is approved
or indicated in monotherapy with said active ingredient.


In another aspect of this invention, there is provided a kit comprising a compound of
formula (I) or (II) and a proton pump inhibitor. The kit is useful as a combined preparation
20 for simultaneous, separate or sequential use in the treatment of patients suffering from
gastrointestinal disorders.

In yet a further aspect of the present invention there is provided a method of making a
pharmaceutical composition comprising a compound of formula (I) or (II) substantially as
25 described herein before and a proton pump inhibitor, comprising mixing said compound
and said proton pump inhibitor with a pharmaceutically acceptable carrier or diluent.

The term "hydrocarbyl" is used herein to refer to monovalent groups consisting of carbon
and hydrogen. Hydrocarbyl groups thus include alkyl, alkenyl and alkynyl groups (in both
30 straight and branched chain forms), cycloalkyl (including polycycloalkyl), cycloalkenyl and
aryl groups, and combinations of the foregoing, such as alkylcycloalkyl,

alkylpolycycloalkyl, alkylaryl, alkenylaryl, alkynylaryl, cycloalkylaryl and cycloalkenylaryl groups.

Where reference is made to a carbon atom of a hydrocarbyl group being replaced by a N, O
5 or S atom, what is intended is that



or that $\text{---CH}_2\text{---}$ is replaced by ---O--- or ---S--- .

A "carbocyclic" group, as the term is used herein, comprises one or more closed chains or
10 rings, which consist entirely of carbon atoms. Carbocyclic groups thus include aryl groups (such as phenyl, naphthyl, indanyl, fluorenyl, (1,2,3,4)-tetrahydronaphthyl, indenyl and isoindenyl, and substituted derivatives thereof), and also alicyclic groups. The term "alicyclic group" refers to a carbocyclic group which does not contain an aromatic ring, and thus includes groups such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl,
15 cyclohexenyl, adamantyl, norbornyl, bicyclo[2.2.2]octyl, norbornenyl and bicyclo[2.2.2]octenyl, and also groups (such as adamantanemethyl and methylcyclohexyl) which contain both alkyl or alkenyl groups in addition to cycloalkyl or cycloalkenyl moieties.

20 The term "aryl" is used herein to refer to an aromatic group, such as phenyl or naphthyl, or a heteroaromatic group, such as pyridyl, pyrrolyl, or furanyl.

The term "alkyl" is used herein to refer to both straight and branched chain forms.

25 When reference is made herein to a substituted carbocyclic group (such as substituted phenyl) or a substituted heterocyclic group, the substituents are preferably from 1 to 3 in number and selected from C_1 to C_6 alkyl, C_1 to C_6 alkoxy, thio, C_1 to C_6 alkylthio, carboxy, carboxy(C_1 to C_6)alkyl, formyl, C_1 to C_6 alkylcarbonyl, C_1 to C_6 alkylcarbonylalkoxy, nitro, trihalomethyl, hydroxy, C_1 to C_6 alkylhydroxy, hydroxy(C_1 to C_6)alkyl, amino, C_1 to C_6
30 alkylamino, di(C_1 to C_6 alkyl)amino, aminocarboxy, C_1 to C_6 alkylaminocarboxy, di(C_1 to C_6 alkyl)aminocarboxy, aminocarboxy(C_1 to C_6)alkyl, C_1 to C_6 alkylaminocarboxy(C_1 to

C₆)alkyl, di(C₁ to C₆ alkyl)aminocarboxy(C₁ to C₆)alkyl, C₁ to C₆ alkylcarbonylamino, C₅ to C₈ cycloalkyl, C₅ to C₈ cycloalkyl(C₁ to C₆)alkyl, C₁ to C₆ alkylcarbonyl(C₁ to C₆ alkyl)amino, aryl, aryl(C₁ to C₆)alkyl, (C₁ to C₆ alkyl)aryl, halo, C₁ to C₆ alkylhalo, sulphamoyl, tetrazolyl and cyano.

5

Most usually, substituents will be selected from C₁ to C₆ alkyl, C₁ to C₆ alkoxy, thio, C₁ to C₆ alkylthio, carboxy, carboxy(C₁ to C₆)alkyl, formyl, C₁ to C₆ alkylcarbonyl, C₁ to C₆ alkylcarbonylalkoxy, nitro, trihalomethyl, hydroxy, C₁ to C₆ alkylhydroxy, hydroxy(C₁ to C₆)alkyl, amino, C₁ to C₆ alkylamino, di(C₁ to C₆ alkyl)amino, aminocarboxy, C₁ to C₆ alkylaminocarboxy, di(C₁ to C₆ alkyl)aminocarboxy, aminocarboxy(C₁ to C₆)alkyl, C₁ to C₆ alkylaminocarboxy(C₁ to C₆)alkyl, di(C₁ to C₆ alkyl)aminocarboxy(C₁ to C₆)alkyl, C₁ to C₆ alkylcarbonylamino, C₅ to C₈ cycloalkyl, C₅ to C₈ cycloalkyl(C₁ to C₆)alkyl, C₁ to C₆ alkylcarbonyl(C₁ to C₆ alkyl)amino, halo, C₁ to C₆ alkylhalo, sulphamoyl, tetrazolyl and cyano.

15

The term "halogen" is used herein to refer to any of fluorine, chlorine, bromine and iodine. Most usually, however, halogen substituents in the compounds of the invention are chlorine and fluorine substituents.

- 20 The term "suitably protected" used herein refers to the use of any suitable protecting group to protect a functional group. Such protecting groups are denoted as PG, PG¹, PG², PG³ etc. as shown in the Reaction Schemes above. Suitable protecting groups will be readily apparent to the skilled person and may be found in, for example, Kocienski, *Protecting Groups*, Thieme, New York, 1994. For example, in the case of hydroxyl groups, suitable
- 25 protecting groups may include esters, ethers (e.g. silyl ethers or alkyl ethers) or acetals. Some specific examples of typical hydroxyl protecting groups are allyl, Aloc, benzyl, BOM, t-butyl, trityl, TBS, TBDPS, TES, TMS, TIPS, PMB, MEM, MOM, MTM, and THP. In the case of nitrogen atoms, suitable protecting groups may include Boc, Aloc, Troc, benzyl, allyl, Fmoc or silyl. In the case of carboxylic acids, suitable protecting groups
- 30 may include esters (e.g. benzyl, allyl, methyl or ethyl esters).

The term "activated" used herein in connection with carboxylic acids refers to any activated derivative of a carboxylic acid. Methods of activating carboxylic acids will be known to the skilled artisan and may include activation using EDC, CDI or DCC (optionally in the presence of nucleophilic catalysts such as DMAP), conversion to an acid
5 halide such as an acid chloride (e.g. using SOCl_2 or oxalyl chloride) or conversion to an activated ester (such as a phenyl or pentafluorophenyl ester).

The invention is now further illustrated by means of the following Examples. All reactions were performed under an atmosphere of dry argon unless otherwise stated.
10 Dichloromethane (DCM) was freshly distilled from calcium hydride. Anhydrous tetrahydrofuran (THF) and N,N-dimethylformamide (DMF) were used.

Example 1 (Reaction Scheme 1)

15 **Step a.** *2-[2-(Adamantan-1-yloxy)-acetyl]-3-dimethylamino-acrylic acid benzyl ester.*

A solution of 4-(adamantan-1-yloxy)-3-oxo-butyric acid benzyl ester (1.133 g, 3.3 mmol) and DMF dimethylacetal (0.56 mL, 3.96 mmol) in DMF (20 mL) was heated at 100°C for 3 h. The mixture was diluted with H_2O (80 mL) and extracted with EtOAc (2×30 mL).
20 The combined extracts were washed with brine (2×30 mL), and dried (MgSO_4). Filtration and evaporation of the solvent afforded the product, (0.647 g, 49%).

^1H NMR. (CDCl_3) δ 7.67 (1H, s), 7.43-7.27 (5H, m), 5.20 (2H, s), 4.34 (2H, s), 3.07-2.89 (6H, b.m), 2.10 (3H, br.s), 1.70 (6H, s), 1.58 (6H, s).

25

Step b. *3-(Adamantan-1-yloxymethyl)-1-o-tolyl-1H-pyrazole-4-carboxylic acid benzyl ester.*

NEt₃ (0.2 mL, 1.43 mmol) was added to a mixture of 2-[2-(adamantan-1-yloxy)-acetyl]-3-dimethylamino-acrylic acid benzyl ester (0.539 g, 1.4 mmol), and o-tolylhydrazine hydrochloride (0.222 g, 1.4 mmol) in MeCN (8 mL) at 0°C. The mixture was allowed to warm to room temperature and stirred for 17 h. The precipitated solid was filtered off,
5 washed with MeCN and dried to afford the product, (0.359 g, 56%).

¹H NMR. (CDCl₃) δ 8.10 (1H, s) 7.48-7.27 (9H, m), 5.33 (2H, s), 4.55 (2H, s) 2.07 (3H, s), 2.02 (3H, br.s), 1.58-1.43 (12H, m).

10 **Step c.** *3-(Adamantan-1-yloxymethyl)-1-o-tolyl-1H-pyrazole-4-carboxylic acid.*

A solution of 3-(adamantan-1-yloxymethyl)-1-o-tolyl-1H-pyrazole-4-carboxylic acid benzyl ester (0.359 g, 0.79 mmol) in MeOH-THF (1:1 / 24 mL) was hydrogenated over 10% palladium on charcoal (50 mg) for 19h. The reaction mixture was filtered through a pad of Celite and the filtrate evaporated to afford the product, (0.29 g, 100%).

15

¹H NMR (DMSO-*d*₆) δ 7.97 (1H, s), 7.46-7.30 (4H, m), 4.53 (2H, s), 1.96 (6H, s), 1.52-1.36 (12H, m).

20 **Step d.** *3-[[3-(Adamantan-1-yloxymethyl)-1-o-tolyl-1H-pyrazole-4-carbonyl]-amino]-benzoic acid benzyl ester.*

Oxalyl chloride (100 µl, 1.15 mmol) was added to a solution of 3-(adamantan-1-yloxymethyl)-1-o-tolyl-1H-pyrazole-4-carboxylic acid (248 mg, 0.56 mmol) and DMF (5 drops) in CH₂Cl₂ (7.5 mL). The reaction mixture was stirred until evolution of carbon dioxide ceased then the solution was evaporated *in vacuo*. The acid chloride obtained was
25 dissolved in THF (5 mL) and added dropwise to a solution of 3-amino-benzoic benzyl ester (146 mg, 0.64 mmol) and *i*-PrNEt₂ (120 µl, 0.69 mmol) in THF (12 mL) and stirred at room temperature for 16h. The solution was diluted with EtOAc (20 mL) washed with 1N HCl (20 mL), saturated NaHCO₃ solution (20mL), brine (20mL), and dried (MgSO₄). Filtration and evaporation afforded the product as a red oil (450 mg, 100%).

30

¹H NMR (CDCl₃) δ 9.78 (1H, s), 8.20 (1H, s), 8.15 (1H, d), 8.10 (1H, s), 7.80 (1H, d), 7.47-7.29 (9H, m), 5.38 (2H, s), 4.36 (2H, s), 2.09 (6H, 2 x s), 1.68 (6H, s), 1.58 (6H, m).

Step e. *3-[[3-(Adamantan-1-yloxymethyl)-1-o-tolyl-1H-pyrazole-4-carbonyl]-amino]-benzoic acid.*

A solution of 3-[[3-(adamantan-1-yloxymethyl)-1-o-tolyl-1H-pyrazole-4-carbonyl]-amino]-benzoic acid benzyl ester (450 mg, 0.8 mmol) in MeOH-THF (10:1 / 22 mL) was hydrogenated over 10% palladium on charcoal (69 mg) for 20 h. The reaction mixture was filtered through a pad of Celite and the filtrate evaporated. The residue was purified by flash column chromatography on silica gel by elution with CH₂Cl₂ /MeOH /acetic acid (9:1:0.1) to afford the product, (290mg, 75%).

10

¹H NMR (DMSO-*d*₆) δ 10.08 (1H, s), 8.33 (1H, s), 8.32 (1H, s), 8.00 (1H, d), 7.66 (1H, d), 7.49-7.33 (5H, m), 4.61 (2H, s), 1.99 (3H, s), 1.96 (3H, s), 1.52-1.38 (12H, m).

The compound was further characterised as the N-methyl-D-glucamine salt. Found: C 60.51, H 7.51, N 7.75%; C₃₆H₄₈N₄O₉•2 H₂O Requires: C 60.32, H 7.31, N 7.82%.

15

Example 2 (Reaction Scheme 2)

Step a. *2-[2-(Adamantan-1-yloxy)-3-acetyl]-3-phenyl-acrylic acid benzyl ester.*

A solution of 4-(adamantan-1-yloxy)-3-oxo-butyric acid benzyl ester (1.138 g, 3.3 mmol), benzaldehyde (0.67 mL, 6.6 mmol) and ethylenediamine diacetate (124 mg, 0.69 mmol) in DMF (15 mL) was stirred at room temperature for 3 h. Pyridine (3 mL) was added and the mixture heated at 70°C for 17 h. The reaction mixture was diluted with 5% KHSO₄ (50 mL) and extracted with EtOAc (2 × 40 mL). The combined extracts were washed with brine (50 mL), and dried (MgSO₄). Filtration and evaporation of the solvent gave the crude product which was a mixture of E and Z isomers and which was purified by flash column chromatography on silica gel, by elution with CH₂Cl₂ /EtOAc (50:1), (729mg, 51%).

25

¹H NMR (CDCl₃) δ (E- isomer) 7.80 (1H, s), 7.43-7.33 (10H, m) 5.28 (2H, s), 4.20 (2H, s), 2.06 (3H, br.s), 1.60-1.48 (12H, m): δ (Z-isomer) 7.80 (1H, s), 7.39-7.27 (10H, m), 5.28 (2H, s), 4.34 (2H, s), 2.14 (3H, br.s), 1.73-1.52 (12H, m).

30

Step b. *3-(Adamantan-1-yloxymethyl)-3-phenyl-1-o-tolyl-1H-pyrazole-4-carboxylic acid benzyl ester.*

A solution of a mixture of E- and Z- isomers of 2-[2-(adamantan-1-yloxy)-3-acetyl]-3-phenyl-acrylic acid benzyl ester (645 mg, 1.5 mmol), *i*-PrNEt₂ (280 µL, 1.6 mmol), o-
5 tolylhydrazine hydrochloride (236 mg, 1.5 mmol) and 1,8-diazabicyclo[5.4.0]undec-7-ene (225 µL, 1.5 mmol) in DMF (8 mL) was heated at 70°C for 16 h. The mixture was partitioned between 5% KHSO₄ (60 mL) and EtOAc (60mL). The organic phase was separated, washed with brine (2 × 60 mL) and dried (MgSO₄). Filtration and evaporation
gave the crude product which was purified by flash column chromatography, by elution
10 with CH₂Cl₂ /EtOAc (40:1). The product was obtained as a yellow oil, (371mg, 46%).

¹H NMR (CDCl₃) δ 7.65 (2H, m), 7.39-7.26 (12H, m), 5.26 (2H, s), 4.54 (2H, s), 2.16 (3H, s), 2.02 (3H, br.s), 1.59-1.43 (12H, m).

15 **Step c.** *3-[[3-(Adamantan-1-yloxymethyl)-3-phenyl-1-o-tolyl-1H-pyrazole-4-carbonyl]-amino]-benzoic acid.*

The title compound was prepared by the same method used to prepare example 1, except that 3-(adamantan-1-yloxymethyl)-3-phenyl-1-o-tolyl-1H-pyrazole-4-carboxylic acid benzyl ester was used in step c in place of 3-(adamantan-1-yloxymethyl)-1-o-tolyl-1H-
20 pyrazole-4-carboxylic acid benzyl ester.

¹H NMR (DMSO-*d*₆) δ 12.87 (1H, br.s), 10.29 (1H, s), 8.28 (1H, s), 7.84 (1H, d), 7.68 (1H, 3H, m), 7.47- 7.33 (8H, m), 4.38 (2H, s), 3.12 (3H, s), 1.93 (3H, br.s), 1.50-1.36 (12H, m).

25 The compound was further characterised as the N-methyl-D-glucamine salt. Found: C 60.47, H 7.15, N 6.95%; C₄₂H₅₂N₄O₉•1.5 H₂O Requires: C 64.35, H 7.07, N 7.15%.

Example 3 (Reaction Scheme 3)

30 **Step a.** *5-(Adamantan-1-yloxymethyl)-2-o-tolyl-2H-pyrazol-3-ol*

A mixture of 4-(adamantan-1-yloxy)-3-oxo-butyric acid benzyl ester (334 mg, 0.98 mmol), o-tolyl hydrazine hydrochloride (163 mg, 1.0 mmol), and NEt₃ (140 µL, 1.0 mmol) in

EtOH (6 mL) was stirred at room temperature for 1 hr. The mixture was evaporated and the residue partitioned between EtOAc (40 mL) and H₂O (40 mL). The organic phase was separated, washed with brine (20 mL), and dried (MgSO₄). Filtration and evaporation of the solvent gave the crude hydrazone. 1,8-Diazabicyclo[5.4.0]undec-7-ene (150 µL, 1.0 mmol) was added to a solution of the hydrazone in DMF (4 mL), and the solution stirred at room temperature for 17 h. The mixture was partitioned between 5% KHSO₄ (60 mL) and EtOAc (60 mL). The organic layer was washed with brine (2 × 30 mL), and dried (MgSO₄). Filtration and evaporation of the solvent gave the product as a solid which was triturated with ether, isolated by filtration and dried (170 mg, 51%).

10

¹H NMR (DMSO-*d*₆) δ 10.44 (1H, s), 7.30 (4H, m), 5.40 (1H, s), 4.27 (2H, s), 2.10 (6H, 2 × s), 1.74 (6H, s), 1.58 (6H, s).

Step b. 3-*[[3-(Adamantan-1-yloxymethyl)-5-hydroxy-1-o-tolyl-1H-pyrazole-4-carbonyl]-amino]-benzoic acid ethyl ester.*

A solution of 5-(adamantan-1-yloxymethyl)-2-o-tolyl-2H-pyrazol-3-ol (160 mg, 0.47 mmol) in anhydrous THF (4 mL) was added to a suspension of sodium hydride (26 mg of 60% dispersion in oil, 0.65 mmol) in THF (2 mL) under a dry argon atmosphere at room temperature. The mixture was stirred for 30 min. and a solution of 3-ethoxycarbonylphenyl isocyanate (99 mg, 0.5 mmol) in THF (1 mL) added. Stirring was continued for 1 h and the mixture partitioned between 5% KHSO₄ (40 mL) and EtOAc (40 mL). The organic phase was washed with brine (30 mL), and dried (MgSO₄). Filtration and evaporation of the solvent afforded the product as a red solid, (250 mg, 100%).

¹H NMR (CDCl₃) δ 10.45 (1H, s), 8.16 (1H, s), 8.03 (1H, d), 7.83 (1H, d), 7.44 (1H, t), 7.28 (4H, m), 4.81 (2H, s), 4.38 (2H, q), 2.22 (3H, s), 2.15 (3H, s), 1.94 (6H, s), 1.65 (6H, s) 1.40 (3H, t).

Step c. 3-*[[3-(Adamantan-1-yloxymethyl)-5-hydroxy-1-o-tolyl-1H-pyrazole-4-carbonyl]-amino]-benzoic acid.*

A solution of 3-*[[3-(adamantan-1-yloxymethyl)-5-hydroxy-1-o-tolyl-1H-pyrazole-4-carbonyl]-amino]-benzoic acid ethyl ester* (249 mg, 0.47 mmol) and potassium hydroxide

(208 mg, 3.7 mmol) in EtOH-H₂O (4:1/ 15 mL) was refluxed for 1 h. On cooling to room temperature the solution was acidified with 5% KHSO₄. The precipitated solid was filtered off, washed with H₂O and dried to afford the product, (201 mg, 85%).

- 5 ¹NMR (DMSO-*d*₆) δ 13.30 (1H, br.s), 12.90 (1H, s), 10.77 (1H, s), 8.28 (1H, s), 7.72 (1H, d), 7.60 (1H, d), 7.45-7.31 (5H, m), 4.91 (2H, s), 2.19 (3H, s) 2.12 (3H, br.s), 1.80 (6H, s), 1.57 (6H, s).

The compound was further characterised as the N-methyl-D-glucamine salt. Found: C
10 59.19, H 7.29, N 7.51%; C₃₆H₄₈N₄O₁₀•2 H₂O Requires: C 59.00, H 7.15, N 7.65%.

Example 4 (Reaction Scheme 3)

Step a. 5-(Adamantan-1-yloxymethyl)-2-cyclohexyl-2H-pyrazol-3-ol

- 15 NEt₃ (800 µl, 5.7 mmol) was added to a solution of 4-(adamantan-1-yloxy)-3-oxo-butyric acid benzyl ester (1.91 g, 5.6 mmol), and cyclohexyl hydrazine hydrochloride (877 mg, 5.82 mmol) in EtOH (30 mL) at 0°C. The mixture was allowed to warm to room temperature and stirred for 4 h. The precipitated white solid was filtered and dried to afford the product as a white solid, (1.23g, 66%).

20

¹H NMR (DMSO-*d*₆) δ 5.31 (1H, s), 4.24 (2H, s), 4.01 (1H, m), 2.18 (3H, br.s), 1.85-1.58 (19H, m), 1.40-1.20 (3H m).

Step b. 3-[[3-(Adamantan-1-yloxymethyl)-1-cyclohexyl-5-hydroxyl-1H-pyrazole-4- 25 carbonyl]-amino]-benzoic acid.

The compound was prepared by the same procedure used in the preparation of Example 3, except that 5-(adamantan-1-yloxymethyl)-2-cyclohexyl-2H-pyrazol-3-ol was used in step b in place of 5-(adamantan-1-yloxymethyl)-2-o-tolyl-2H-pyrazol-3-ol.

- 30 ¹H NMR (CDCl₃) δ 10.75 (1H, br.s), 10.34 (1H, s), 8.25 (1H, s), 7.98 (1H, d), 7.86 (1H, d), 7.47 (1H, t), 4.73 (2H, s), 4.12 (1H, m), 2.20 (1H, br. s), 1.92 (13H, m), 1.66 (6H, m), 1.43-1.19 (3H, m).

The compound was further characterised as the N-methyl-D-glucamine salt. Found: C 58.12, H 7.87, N 7.58%; $C_{35}H_{52}N_4O_{10} \cdot 2 H_2O$ Requires: C 58.00, H 7.79, N 7.73%.

5 Example 5 (Reaction Scheme 4)

Step a. 3-(Adamantan-1-yloxymethyl)-1-cyclohexyl-5-methoxy-1H-pyrazole

5-(Adamantan-1-yloxymethyl)-2-cyclohexyl-2H-pyrazol-3-ol (0.29 g, 0.88 mmol) and sodium hydride (50 mg, 1.25 mmol of a 40% suspension in mineral oil) were stirred in THF-DMF (1:1) (10 mL) at 0 °C for 0.5 h, then allowed to warm to room temperature. Dimethyl sulphate (0.1 mL, 1 mmol) was added and the mixture stirred overnight. After concentration in vacuo, the mixture was poured into water (10 mL) and extracted with EtOAc (20 mL). The organic layer was separated washed with water (2 x 10 mL) and dried (MgSO₄). Filtration and evaporation of the solvent afforded a yellow oil, which was purified by chromatography with EtOAc-hexane (1:1) as eluant. (0.24 g, 81%).

¹H NMR (CDCl₃) δ 5.54 (1H, s), 4.40 (2H, s), 3.96 (1H, m), 3.84 (3H, s), 2.15 (3H, br.s), 1.97-1.82 (12H, m), 1.67-1.63 (7H, m), 1.37-1.24 (m, 3H).

20 Step b. 3-(Adamantan-1-yloxymethyl)-4-bromo-1-cyclohexyl-5-methoxy-1H-pyrazole

3-(Adamantan-1-yloxymethyl)-1-cyclohexyl-5-methoxy-1H-pyrazole (0.24 g, 0.7 mmol), azobisisobutyronitrile (2 mg) and N-bromosuccinimide (0.14 g, 0.8 mmol) were stirred in CCl₄ (10 mL) for 1h under illumination (100 W light source). After filtration, the filtrate was concentrated *in vacuo* to afford a yellow oil. (0.3 g, 100%)

25

¹H NMR (CDCl₃) δ 4.36 (2H, s), 4.03-4.01 (4H, m), 2.14 (3H, br.s), 1.86-1.61 (19H, m), 1.33-1.26 (3H, m).

Step c. 3-[[3-(Adamantan-1-yloxymethyl)-1-cyclohexyl-5-methoxy-1H-pyrazole-4-carbonyl]-amino}-benzoic acid ethyl ester

30

The compound was prepared by treating 3-(adamantan-1-yloxymethyl)-4-bromo-1-cyclohexyl-5-methoxy-1H-pyrazole with n-butyl lithium and 3-ethoxycarbonylphenyl isocyanate as outlined in Reaction Scheme 4.

5 ¹H NMR (CDCl₃) δ 10.22 (1H, s), 8.09 (2H, d), 7.77 (1H, d), 7.42 (1H, t), 4.66 (2H, s), 4.38 (2H, q), 4.12 (4H, m), 2.18 (3H, br.s), 1.91-1.64 (22H, m), 1.25 (6H, t).

Step d. 3-{[3-(Adamantan-1-yloxymethyl)-1-cyclohexyl-5-methoxy-1H-pyrazole-4-carbonyl]-amino}-benzoic acid

10 The compound was prepared by the same procedure used in the preparation of Example 3, except that 3-{[3-(adamantan-1-yloxymethyl)-1-cyclohexyl-5-methoxy-1H-pyrazole-4-carbonyl]-amino}-benzoic acid ethyl ester was used in step c in place of 3-{[3-(adamantan-1-yloxymethyl)-5-hydroxy 1-o-tolyl-1H-pyrazole-4-carbonyl]-amino}-benzoic acid ethyl ester.

15

¹H NMR (DMSO-d₆) δ 12.92 (1H, br.s), 10.07 (1H, s), 8.20 (1H, s), 7.88 (1H, d), 7.63 (1H, s), 7.46 (1H, t), 4.54 (2H, s), 4.11 (1H, m), 4.00 (3H, s), 2.05 (3H, br.s), 1.79-1.21 (22H, m). MS (EI, MeOH), 507 (M⁺, 3%), 372 (M⁺, -Ad, 12%) calcd. 507 for C₂₉H₃₇N₃O₅.

20 **Example 6 (Reaction Scheme 5)**

Step a. 5-(Adamantan-1-yloxymethyl)-4-bromo-2-cyclohexyl-1,2-dihydro-pyrazol-3-one

A CCl₄ solution (10 mL) of 5-(adamantan-1-yloxymethyl)-2-cyclohexyl-2H-pyrazol-3-ol (0.11 g, 0.33 mmol), azobisisobutyronitrile (2 mg) and N-bromosuccinimide (70 mg, 0.4 mmol) was irradiated by a 100 W light source for 0.3 h at rt. After filtration to the filtrate was concentrated *in vacuo* to afford a crude product as a purple solid, which was purified by chromatography with EtOAc-DCM (1:4) as eluant. The product was obtained as a white glassy solid (65mg, 48%).

30 ¹H NMR (CDCl₃) δ 4.89 (1H, s), 4.18 (2H, m), 3.96 (1H, m), 2.19 (3H, br.s), 1.84-1.26 (22H, m).

¹H NMR (DMSO-d₆) δ 10.9 (1H, br.s), 4.18 (2H, s), 4.04 (1H, m), 2.09 (3H, br.s), 1.79-1.30 (22H, m).

Step b. *5-(Adamantan-1-yloxymethyl)-4-bromo-2-cyclohexyl-1-methyl-1,2-dihydro-pyrazol-3-one*

The compound was prepared by treating 5-(adamantan-1-yloxymethyl)-4-bromo-2-cyclohexyl-1,2-dihydro-pyrazol-3-one with sodium hydride and methyl iodide as outlined in Reaction Scheme 5.

¹H NMR (CDCl₃) δ 11.01 (1H, s), 8.15-8.08 (2H, m), 7.73 (1H, d), 7.35 (1H, t), 5.15 (2H, s), 4.36 (2H, q), 4.08 (1H, m), 3.71 (3H, s), 2.25 (5H, m), 1.97-1.64 (17H, m), 1.40-1.27 (6H, m).

Step c. *3-{{5-(Adamantan-1-yloxymethyl)-2-cyclohexyl-1-methyl-3-oxo-2,3-dihydro-1H-pyrazole-4-carbonyl]-amino}-benzoic acid*

The compound was prepared by the same procedure used in the preparation of Example 5, except that 3-{{5-(adamantan-1-yloxymethyl)-2-cyclohexyl-1-methyl-3-oxo-2,3-dihydro-1H-pyrazole-4-carbonyl]-amino}-benzoic acid ethyl ester was used in step c in place of 3-{{3-(adamantan-1-yloxymethyl)-1-cyclohexyl-5-methoxy-1H-pyrazole-4-carbonyl]-

amino}-benzoic acid ethyl ester

¹H NMR (DMSO-d₆) δ 12.92 (1H, br.s), 11.20 (1H, s), 8.27 (1H, s), 7.72 (1H, d), 7.61 (1H, s), 7.42 (1H, t), 5.00 (2H, s), 4.13 (1H, m), 3.67 (3H, s), 2.05 (5H, m), 1.79-1.18 (20H, m). MS (EI, MeOH), 507 (M⁺, 1%), 372 (M⁺, -Ad, 20%) calcd. 507 for C₂₉H₃₇N₃O₅.

25

The compounds of the examples were tested for gastrin (CCK₂) antagonist activity in an immature rat stomach assay. The procedure was as follows:

The oesophagus of immature rats (33-50 g, ca. 21 days old) was ligated at the level of the cardiac sphincter and the duodenal sphincter was cannulated. The stomach was excised and flushed with ca. 1 ml of unbuffered physiological saline solution. The fundus was punctured and cannulated. A further 4-5 ml of unbuffered solution was flushed through the

stomach to ensure the preparation was not leaking. The stomach was lowered into a jacketed organ bath containing 40 ml of buffered solution containing 3×10^{-8} M 5-methylfurmethide, maintained at 37° and gassed vigorously with 95% O_2 / 5% CO_2 . The stomach was continuously perfused at a rate of 1 ml min^{-1} with unbuffered solution gassed with 100% O_2 with the perfusate passing over an internally referenced pH-electrode fixed 12 cm above the stomach.

After 120 min of stabilisation the drugs were added directly to the serosal solution in the organ bath and after a further 60 min cumulative pentagastrin dose-response curves were started. Changes in acid secretion were monitored and the curves analysed according to Black *et al.*, *Br. J. Pharmacol.*, 1985, **86**, 581.

The results obtained at gastrin (CCK_2) receptors are set out in Table 1.

15

Table 1

Example	Rat Stomach pK_B
2	5.66 ± 0.24
3	7.72 ± 0.20
4	7.51 ± 0.30
5	6.77 ± 0.38
6	6.57 ± 0.30

It is found that the compositions and products of the present invention comprising a compound of formula (I) and a proton pump inhibitor reduce hyperplasia, associated with administration of proton pump inhibitors. This was measured according to the following experimental protocol.

20

Animals and treatment:

40 male SPF Wistar rats (200 g) were divided into 4 treatment groups and 2 strata. The treatment of the 20 rats in the second stratum started 2 weeks after the treatment of the first stratum. The design of the study was completely randomised double blind with individual
5 blinding; all rats were placed in a separate cage. Animals had continuous access to water and food.

Animals were treated once daily during 14 days:

- Control group: 1 ml gastrin test drug vehicle + 1 ml p.o.(gavage) 0,25% Methocel
10 (Dow Corning)
- PPI group: 1 ml gastrin test drug vehicle +1 ml p.o.(gavage) 25 mg/kg Rabeprazole
in 0.25% Methocel.
- GRA group: 1 ml gastrin test drug + 1 ml p.o. (gavage) 0,25% Methocel
- GRA-PPI group: 1 ml gastrin test drug + 1 ml p.o.(gavage) 25 mg/kg Rabeprazole in
15 0.25% Methocel.

Gastrin test drug made up to an appropriate dose in physiologically compatible solvent.

Preparation of tissue:

- 20 After removal of the fundus, the stomach were rinsed with phosphate buffered saline prior to fixation with 4% formalin in Millonig buffer. After 4 hours immersion in fixative solutions at room temperature, tissue was rinsed in phosphate buffered saline (PBS), dehydrated and embedded in paraffin using the Leitz paraffin embedding station (Leitz TP 1050; Germany) dehydration module and paraffin embedding module (Leitz EG 1160;
25 Germany).

Cross sections (3 μ m thick) of the oxyntic part of the stomach were made at 3 levels, each separated by a distance of 400 μ m.

30 Immunostaining

The following indirect immunofluorescence labeling method was used:

- removal of paraffin and rehydration of the sections followed by a blocking step

- primary antibodies: polyclonal guinea pig anti-histidine decarboxylase, 1/2000 (from Euro-Diagnostica) and monoclonal mouse anti PCNA 1/2500 (Clone PC10 from Sigma). All antibodies were diluted in a 0.2% BSA solution. Sections were incubated overnight at 4°C and then washed with a BSA solution.
- 5 - secondary antibodies: goat anti guinea pig coupled to CY5, 1/500 (from Jackson Laboratories) and goat anti-mouse coupled to Cy3, 1/250 (from Jackson Laboratories); incubation for 4 hours at 37°C. After rinsing with BSA and PBS solutions, sections were mounted with slowfade (Molecular Probes Europe BV), and stored at 4°C.

10 Imaging

Fluorescence labelling was observed with an epifluorescence microscope or a Zeiss LSM510 (Carl Zeiss Jena GmbH) confocal microscope.


By using CY5- and CY3-coupled antibodies, the high autofluorescence properties of the oxyntic mucosa were circumvented when sections are illuminated by a 488 nm (FITC channel) light source. Negative controls, by omitting the primary antibodies, and an isotype control staining for PCNA showed complete absence of staining. The specific labelling of PCNA was checked using double staining with TOPRO-3® (Molecular Probes Europe BV), a nuclear stain. Only in the most luminal located epithelial cells, non-specific
15 cytoplasmic labelling was present. In the glandular part of the mucosa, non-specific PCNA-staining was absent.

For determination of the labelling index of ECL cells, at least 80 confocal images per rat were taken from the 3 slides at the 3 different levels. The ratio of double labelled cells
25 (HDC + PCNA) and all HDC labelled cells yielded the labelling index of ECL cells.

Proliferation activity of ECL cells in the PPI group is expected to be increased compared with sham, GRA and GRA-PPI groups (Eissele, R., Patberg, H., Koop, H., Krack, W., Lorenz, W., McKnight, A.T., and Arnold, R. Effect of gastrin receptor blockade on
30 endocrine cells in rats during achlorhydria. *Gastroenterology*, 103, 1596-1601, 1992). Increased proliferation by PPI will be completely blocked by GRA.

Claims

1. A compound of the formula (I), (IIa) or (IIb)



wherein


- 5 n is from 1 to 4;
- R^1 is C_1 to C_{15} hydrocarbyl wherein up to three C atoms may optionally be replaced by N, O and/or S atoms and up to three H atoms may optionally be replaced by halogen atoms;
- R^2 is selected from H, Me, Et, Pr and OH, each R^2 being independently selected from H, Me, Et, Pr and OH when n is greater than 1;
- 10 R^3 (when n is 1) is selected from H, Me, Et and Pr; or (when n is greater than 1) each R^3 is independently selected from H, Me, Et and Pr, or two R^3 groups on neighbouring carbon atoms are linked to form a C_3 to C_6 carbocyclic ring, or two R^3 groups are absent from neighbouring carbon atoms which are linked by a
- 15 double bond; or R^2 and R^3 on the same carbon atom together represent an =O group;
- R^4 is C_1 to C_{15} hydrocarbyl wherein up to two C atoms may optionally be replaced by N, O and/or S atoms and up to three H atoms may optionally be replaced by halogen atoms;
- 20 R^5 is H, OH or C_1 to C_{15} hydrocarbyl wherein up to two C atoms may optionally be replaced by N, O and/or S atoms and up to three H atoms may optionally be replaced by halogen atoms;

R^6 is H or C_1 to C_{15} hydrocarbyl wherein up to two C atoms may optionally be replaced by N, O and/or S atoms and up to three H atoms may optionally be replaced by halogen atoms;


5 Z is $-(NR^7)_a-CO-(NR^8)_b-$ (wherein a is 0 or 1, b is 0 or 1, and R^7 and R^8 are independently selected from H, Me, Et, Pr or Bn),

$-CO-NR^7-CH_2-CO-NR^8-$, $-CO-O-$, $-CH_2-CH_2-$, $-CH=CH-$, $-CH_2-NR^8-$ or a bond;

Q is $-R^9V$, or



10 wherein R^9 is $-CH_2-$; $-CH_2-CH_2-$; or



or R^9 and R^8 , together with the nitrogen atom to which R^8 is attached, form a piperidine or pyrrolidine ring which is substituted by V;

15 V is $-CO-NH-SO_2-Ph$, $-SO_2-NH-CO-Ph$, $-CH_2OH$, or a group of the formula $-R^{10}U$, (wherein U is $-COOH$, tetrazolyl, $-CONHOH-$ or $-SO_3H$; and R^{10} is a bond; C_1 to C_6 hydrocarbylene, optionally substituted by hydroxy, amino or acetamido; $-O-(C_1$ to C_3 alkylene)-; $-SO_2NR^{11}-CHR^{12}-$;

$-CO-NR^{11}-CHR^{12}-$, R^{11} and R^{12} being independently selected from H and methyl; or $-NH-(CO)_c-CH_2-$, c being 0 or 1);

20 T is C_1 to C_6 hydrocarbyl, $-NR^7R^8$ (wherein R^7 and R^8 are as defined above), $-OMe$, $-OH$, $-CH_2OH$, halogen or trihalomethyl;

m is 1 or 2;


p is from 0 to 3; and

q is from 0 to 2, with the proviso that q is 1 or 2 when Z is a bond);


25 or a pharmaceutically acceptable salt thereof.

2. A compound according to claim 1 wherein Q is

35




3. A compound according to claim 2 wherein Q is



- 5 4. A compound according to any preceding claim wherein R^1 is C_1 to C_{12} hydrocarbyl, wherein one C atom may optionally be replaced by N or O and up to three H atoms may optionally be replaced by F, Cl or Br.
5. A compound according to any preceding claim wherein R^1 is C_3 to C_{12} alicyclic; phenyl, pyridyl, phenyl C_1 - C_3 alkyl or pyridyl C_1 - C_3 alkyl (all optionally substituted with OMe, NMe₂, CF₃, Me, F, Cl, Br or I); or C_1 to C_8 alkyl.
- 10 6. A compound according to any preceding claim wherein R^5 is H, OH or C_1 to C_{12} hydrocarbyl wherein one C atom may optionally be replaced by N or O and up to three H atoms may optionally be replaced by F, Cl, or Br.
- 15 7. A compound according to any preceding claim wherein R^5 is H; OH; C_3 to C_{12} alicyclic; O- C_3 to C_{12} alicyclic; phenyl, pyridyl, phenyl C_1 - C_3 alkyl, phenyl C_1 - C_3 alkoxy, pyridyl C_1 - C_3 alkyl, or pyridyl C_1 - C_3 alkoxy (all optionally substituted with OMe, NMe₂, CF₃, Me, F, Cl, Br or I); C_1 to C_8 alkyl; or C_1 to C_8 alkoxy.
- 20 8. A compound according to any preceding claim wherein R^6 is H or C_1 to C_{12} hydrocarbyl wherein one C atom may optionally be replaced by N or O and up to three H atoms may optionally be replaced by F, Cl, or Br.
- 25 9. A compound according to any preceding claim wherein R^6 is H; C_3 to C_{12} alicyclic; phenyl, pyridyl, phenyl C_1 - C_3 alkyl or pyridyl C_1 - C_3 alkyl (all optionally substituted with OMe, NMe₂, CF₃, Me, F, Cl, Br or I); or C_1 to C_8 alkyl.

10. A compound according to any preceding claim wherein Z is $-\text{CO}-\text{NH}-$.
11. A compound according to any preceding claim wherein p is 0 or 1, and q is 0.
- 5 12. A compound according to any preceding claim wherein T is C_1 to C_6 hydrocarbyl or halo.
13. A compound according to any preceding claim wherein V is $-\text{CO}_2\text{H}$,
10 $-\text{CH}_2\text{CO}_2\text{H}$ or tetrazolyl.
14. A compound according to any preceding claim wherein R^2 and R^3 are H, and n is from 1 to 3.
- 15 15. A compound according to any of claims 1 to 13 wherein R^2 and R^3 together form an $=\text{O}$ group, and n is 1.
16. A compound according to claim 14 or claim 15 wherein R^4 is C_3 to C_{12} carbocyclic.
- 20 17. A compound according to claim 16 wherein R^4 is adamantyl, cycloheptyl, cyclohexyl or phenyl.
18. A compound according to claim 14 or claim 15 wherein R^4 is $-\text{NH}-\text{R}^{13}$ or $-\text{OR}^{13}$,
in which R^{13} is C_3 to C_{12} carbocyclic.
- 25 19. A compound according to claim 18 wherein R^{13} is adamantyl, cycloheptyl, cyclohexyl or phenyl.
20. A compound according to claim 1 wherein Z is $-(\text{NR}^7)_a-\text{CO}-(\text{NR}^8)_b-$, $-\text{CO}-\text{NH}-$
30 $\text{CH}_2-\text{CO}-\text{NH}-$ or a bond; Q is

37




V is $-\text{CO}-\text{NH}-\text{SO}_2-\text{Ph}$, $-\text{SO}_2-\text{NH}-\text{CO}-\text{Ph}$, $-\text{OCH}_2\text{COOH}$, tetrazolyl or $-(\text{CH}_2)_s\text{COOH}$, wherein s is from 0 to 2; and T is C_1 to C_6 hydrocarbyl, $-\text{NR}^8\text{R}^8$, $-\text{OMe}$, $-\text{OH}$, $-\text{CH}_2\text{OH}$ or halogen.

5


21. A compound according to claim 1 wherein Z is $-(\text{NR}^7)_a-\text{CO}-(\text{NR}^8)_b-$, Q is $-(\text{CH}_2)_r\text{COOH}$, wherein r is from 1 to 3; and T is C_1 to C_6 hydrocarbyl, $-\text{NR}^7\text{R}^8$, $-\text{OMe}$, $-\text{OH}$, $-\text{CH}_2\text{OH}$ or halogen.

10 22. A compound according to claim 1 wherein $-\text{Z}-\text{Q}$ is



23. A compound which is degraded *in vivo* to yield a compound according to any
15 preceding claim.


24. A method of making a compound of formula (I) wherein R^5 is H including the step of reacting a compound of formula (IV), or a suitably protected derivative thereof, with R^1-NHNH_2 in the presence of a base.




20

25. A method of making a compound of formula (I) wherein R^5 is C_3 to C_{12} alicyclic; phenyl (optionally substituted with OMe , NMe_2 , CF_3 , Me , F , Cl , Br , I or OH); or C_1 to C_8 alkyl, including the step of reacting a compound of formula (VII), or a suitably protected
25 derivative thereof, with R^1-NHNH_2 in the presence of a base.

38



26. A method of making a compound of formula (I) wherein R⁵ is OH or a compound of formula (IIa) or (IIb), including the step of reacting a compound of formula (III), or a suitably protected derivative thereof, with R¹-NHNH₂ in the presence of a base.



27. A pharmaceutical composition comprising a compound of formula (I) or (II) according to any of claims 1 to 23 together with a pharmaceutically acceptable diluent or carrier.

28. A pharmaceutical composition comprising a proton pump inhibitor and a compound of formula (I) or (II) according to any one of claims 1 to 23 together with a pharmaceutically acceptable diluent or carrier.

29. A composition according to claim 28 wherein the proton pump inhibitor is selected from (RS)-rabeprazole, (RS)-omeprazole, lansoprazole, pantoprazole, (R)-omeprazole, (S)-omeprazole, perprazole, (R)-rabeprazole, (S)-rabeprazole, or the alkaline salts thereof.

30. A composition according to claim 28 or 29 wherein the proton pump inhibitor and the compound of formula (I) or (II) are each in an amount producing a therapeutically beneficial effect in patients suffering from gastrointestinal disorders.

31. A composition according to claim 30 wherein said therapeutically beneficial effect is a synergistic effect on the reduction of acid secretion in patients suffering from gastrointestinal disorders, or the prevention of gastrointestinal disorders in said patients, or

the reduction of adverse effects associated with the one of the active ingredients by the other active ingredients.

32. A composition according to any one of claims 28 to 31 wherein the amount of each
5 of the active ingredients is equal to or less than that which is approved or indicated in monotherapy with said active ingredient.

33. A product containing as first active ingredient a compound of formula (I) or (II) and
as second active ingredient a proton pump inhibitor, as a combined preparation for
10 simultaneous, separate or sequential use in the treatment of patients suffering from gastrointestinal disorders.

34. A compound according to any one of claims 1 to 23, a composition according to
any one of claims 28 to 32, or a product according to claim 33 for use in medicine.

15

35. A compound according to any one of claims 1 to 23, a composition according to
any one of claims 27 to 32, or a product according to claim 33 for use in treating
gastrointestinal disorders.

20 36. Use of a compound according to any one of claims 1 to 23, a composition according
to any one of claims 27 to 32 or a product according to claim 33 for the preparation of a
medicament for the treatment of gastrointestinal disorders.

37. Use of a proton pump inhibitor for the preparation of a medicament for the
25 treatment of gastrointestinal disorders, said treatment comprising the simultaneous or
sequential administration of said proton pump inhibitor and a compound of formula (I) or
(II), wherein said proton pump inhibitor enhances the effect of the compound of formula (I)
or (II) on gastrin-related disorders in patients suffering from gastrointestinal disorders.

30 38. Use of a compound of formula (I) or (II) for the preparation of a medicament for the
treatment of gastrointestinal disorders, said treatment comprising the simultaneous or
sequential administration of said proton pump inhibitor and a compound of formula (I) or

(II), wherein said compound of formula (I) or (II) enhances the effect of the proton pump inhibitor on the reduction of acid secretion in patients suffering from gastrointestinal disorders.

5 39. Use of a compound of formula (I) or (II) for the preparation of a medicament for reducing adverse effects associated with administration of proton pump inhibitors in patients suffering from gastrointestinal disorders.

40. Use according to claim 39 wherein the adverse effect is hyperplasia.

10

41. Use of a proton pump inhibitor for the preparation of a medicament for reducing adverse effects associated with administration of a compound of formula (I) or (II) in patients suffering from gastrointestinal disorders.

15 42. Use of a compound according to any one of claims 1 to 23, a composition according to any one of claims 27 to 32 or a product according to claim 33 in the treatment of gastrointestinal disorders.

43. A method of making a pharmaceutical composition according to claim 27
20 comprising mixing a compound of formula (I) or (II) with a pharmaceutically acceptable diluent or carrier.

44. A method of making a pharmaceutical composition according to any one of claims
28 to 32, comprising mixing a compound of formula (I) or (II) and a proton pump inhibitor
25 with a pharmaceutically acceptable diluent or carrier.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 01/01976

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D231/14 C07D231/20 C07D231/22 A61K31/415 A61P1/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EP0-Internal, WPI Data, PAJ, CHEM ABS Data, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4 670 460 A (MARDIN MITHAT ET AL) 2 June 1987 (1987-06-02) column 2, line 1 - line 50 ---	1,27,34
X	DE 23 63 139 A (BAYER AG) 10 July 1975 (1975-07-10) claims 1,3 ---	1,27,34
X	FR 2 773 153 A (RHONE POULENC AGROCHIMIE) 2 July 1999 (1999-07-02) claim 1 ---	1
X	SUCROW, WOLFGANG ET AL: "Enehydrazines, 12. Some products from methyl 1-alkyl-5-hydroxy-3- pyrazolecarboxylates" CHEM. BER. (1976), 109(1), 261-7, XP001007940 * Compounds 11 and 12 * ---	1
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the International filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the International filing date but later than the priority date claimed

- "T" later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the International search

11 July 2001

Date of mailing of the International search report

21.02.01

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Johnson, C

INTERNATIONAL SEARCH REPORT

Inte I Application No
PCT/GB 01/01976

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,Y	WO 00 27823 A (BLACK JAMES WHYTE ;BUCK ILDIKO MARIA (GB); DUNSTONE DAVID JOHN (GB) 18 May 2000 (2000-05-18) cited in the application claims 1,28	1,27,34, 35
Y	--- EP 0 467 614 A (LILLY CO ELI) 22 January 1992 (1992-01-22) claims 1,19	1,27,34, 35
Y	--- US 5 202 344 A (BECKER DANIEL P ET AL) 13 April 1993 (1993-04-13) claims 1,23 -----	1,27,34, 35

INTERNATIONAL SEARCH REPORT

ational application No.
PCT/GB 01/01976

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 23
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1 - 36, 38 - 40, 42 - 44

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 23

Present claim 23 relates to a compound defined by reference to a desirable characteristic or property, namely that it is degraded in vivo to yield a compound of formula I or II. The scope of the claims must be unambiguously clear to the skilled man. Although the general principle of prodrugs is known, in order to determine which of the many various functional groups of the compounds of formula I or II can be protected by which potentially degradable groups to give a compound falling within the scope of claim 23, the skilled man would be required to perform extensive in vivo testing. As the scope of the claim is not clear without such testing, this claim is not considered to comply with the requirements of Article 6 PCT.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-36,38-40,42-44

Compounds of formula (I), (IIa) and (IIb), their preparation, pharmaceutical compositions, first and second medical uses.

2. Claims: 37,41

Second medical uses of proton pump inhibitors.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/GB 01/01976

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 4670460	A	02-06-1987	DE 3308881 A	13-09-1984
			DE 3342628 A	11-07-1985
			AU 2460884 A	13-09-1984
			DK 143684 A	13-09-1984
			EP 0125406 A	21-11-1984
			ES 530432 D	16-04-1985
			ES 8503945 A	01-07-1985
			FI 840954 A	13-09-1984
			GR 81867 A	12-12-1984
			HU 34123 A	28-02-1985
			JP 59175469 A	04-10-1984
			NO 840869 A	13-09-1984
			PT 78239 A,B	01-04-1984
			ZA 8401766 A	31-10-1984
DE 2363139	A	10-07-1975	AR 207444 A	08-10-1976
			AR 204951 A	19-03-1976
			AR 208192 A	09-12-1976
			AR 205271 A	21-04-1976
			AR 205272 A	21-04-1976
			AT 337176 B	10-06-1977
			AT 313274 A	15-10-1976
			AT 347933 B	25-01-1979
			AT 325876 A	15-06-1978
			AT 347934 B	25-01-1979
			AT 325976 A	15-06-1978
			AT 348514 B	26-02-1979
			AT 326076 A	15-07-1978
			AT 348515 B	26-02-1979
			AT 326176 A	15-07-1978
			AU 6817474 A	23-10-1975
			BG 21217 A	20-03-1976
			BG 21857 A	20-09-1976
			BG 21218 A	20-03-1976
			BG 22821 A	20-04-1977
			BG 21405 A	20-05-1976
			CA 1034953 A	18-07-1978
			CH 596172 A	28-02-1978
			CH 605801 A	13-10-1978
			CH 605800 A	13-10-1978
			CH 605797 A	13-10-1978
			CH 603592 A	31-08-1978
			CS 189648 B	30-04-1979
			CS 189649 B	30-04-1979
			CS 189650 B	30-04-1979
			DD 113757 A	20-06-1975
			DK 136066 B	08-08-1977
			EG 11157 A	29-03-1978
			ES 425373 A	01-09-1976
			ES 447063 A	16-06-1977
			ES 447064 A	16-09-1977
			FI 57587 B	30-05-1980
			FR 2226172 A	15-11-1974
			GB 1429141 A	24-03-1976
			HK 70276 A	19-11-1976
			HU 168590 B	28-06-1976
			IE 39212 B	30-08-1978
			IL 44626 A	31-01-1980

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/GB 01/01976

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 2363139 A		JP 934883 C JP 50018466 A JP 53014071 B LU 69872 A NL 7405115 A,B, NO 138530 B PH 14829 A	30-11-1978 26-02-1975 15-05-1978 28-11-1974 21-10-1974 12-05-1978 16-12-1981
FR 2773153 A	02-07-1999	AU 1882099 A WO 9933813 A	19-07-1999 08-07-1999
WO 0027823 A	18-05-2000	AU 1061500 A NO 20012288 A	29-05-2000 02-07-2001
EP 0467614 A	22-01-1992	AU 5529394 A AU 644190 B AU 8040991 A BR 9103042 A CA 2046672 A CN 1058209 A,B CS 9102193 A FI 913431 A HU 58296 A IE 912489 A IL 98785 A JP 4230367 A KR 9605950 B MX 9100164 A NO 912772 A NZ 238912 A PT 98323 A RU 2076100 C US 5300514 A US 5399565 A US 5643926 A ZA 9105372 A	28-04-1994 02-12-1993 23-01-1992 28-04-1992 18-01-1992 29-01-1992 19-02-1992 18-01-1992 28-02-1992 29-01-1992 27-11-1995 19-08-1992 06-05-1996 28-02-1992 20-01-1992 23-12-1992 30-06-1992 27-03-1997 05-04-1994 21-03-1995 01-07-1997 31-03-1993
US 5202344 A	13-04-1993	AT 116638 T AU 9057191 A CA 2097517 A DE 69106532 D DE 69106532 T DK 561941 T EP 0561941 A ES 2067322 T IE 914293 A,B JP 6503827 T PT 99739 A WO 9210476 A US 5314886 A	15-01-1995 08-07-1992 12-06-1992 16-02-1995 08-06-1995 06-03-1995 29-09-1993 16-03-1995 17-06-1992 28-04-1994 30-11-1992 25-06-1992 24-05-1994